

READ ENTIRE LABEL BEFORE USE. SHAKE WELL BEFORE USE.

KEEP CONTAINER CLOSED WHEN NOT IN USE. DO NOT STORE IN DIRECT SUNLIGHT FOR PROLONGED PERIOD OF TIME.

CAUTION:

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. In case of contact with eyes, flush immediately with copious amounts of water for 15 minutes. If irritation persists, consult a physician. Avoid contamination of skin and clothing. Wash with soap and water.

WARRANTY AND CONDITION OF SALE: Manufacturer warrants that this material conforms to the chemical description hereon and is reasonably fit for use as directed. No other warranty is made, either expressed or implied. Buyer and user, therefore, acknowledges and assumes all risk and liability resulting from handling, storage, and use of this material.

Manufactured By: ProTea Botan US, INC.

1039 Autumn Oaks Circle Collierville TN 38017

FOR FLORIDA:

NET CONTENTS: 2.5 USG / 9.45 L 20.95 lbs / 9.52 Kg

We recommend that you follow the Golf Course BMP's at: http://www.flaes.org/pdf/Golf%20Course%20BMP's.pdf

COLONY SOIL INNOCULANT PROFESSIONAL TURF USE

DESCRIPTION:

COLONY is a concentrated proprietary blend of select soil bacteria that is designed for use on all intensively managed turfgrass. Program use of this product enables the soil to maintain a healthy level of essential beneficial organisms that are often compromised in the commercial turf environment.

COMPATIBILITY:

COLONY can be tank mixed with any nutrient solution that does not contain copper. Can be tank mixed with Organic Matters, Penetrar, Retenador, and CaLpHixx.

✤ DO NOT TANK MIX WITH PESTICIDES.

APPLICATION:

Apply as a complete cover spray to target area. Water in immediately.

GENERAL RECOMMENDATIONS:

Apply 4-6 fl. oz. per 1000 f^2 (120-180 ml/100 m²). Repeat on a monthly basis as long as temperatures are above 65°f, (18°C).

- For Black Algae affected areas: Apply 12-15 fl. oz. per 1000 f², (360-450 ml/100 m²). Maintain light surface moisture and repeat as needed. (CORRECT INFILTRATION PROBLEMS).
- Laying Sod: Apply at 4-6 fl. oz. per 1000 f², (120-180 ml/100 m²), in conjunction with Organic Matters immediately prior to laying sod